

Rapid Molecular Identification of *Tetraselmis* Using Enzymatic Digestion of the 18S *rDNA* Gene

Janeth Galarza^{1,*}, Kevin Crespín¹, Carolina Tufiño²

¹Biology Research Center, Faculty of Marine Sciences, Santa Elena Peninsula State University, La Libertad, Ecuador

²Laboratory of Marine Physiology and Genetics, Center for Advanced Studies in Arid Zones, Coquimbo, Chile

Email address:

jgalarza@upse.edu.ec (J. Galarza)

*Corresponding author

To cite this article:

Janeth Galarza, Kevin Crespín, Carolina Tufiño. Rapid Molecular Identification of *Tetraselmis* Using Enzymatic Digestion of the 18S *rDNA* Gene. *International Journal of Genetics and Genomics*. Vol. 8, No. 3, 2020, pp. 114-119. doi: 10.11648/j.ijgg.20200803.14

Received: July 18, 2020; Accepted: August 17, 2020; Published: September 24, 2020

Abstract: *Tetraselmis* is a genus of quadriflagellated single-celled green algae belonging to the Phylum Chlorophyta, commonly used in aquaculture with very promising biotechnological potential. The varied morphological characteristics, in some cases, have led to confusion in taxonomic identification. To solve this problem, new techniques based on molecular markers and restriction enzymes can ensure the identification of microalgae without sequencing. This study aimed to compare *in silico* modeling with an experimental restriction pattern based on the 18S *rDNA* gene for the identification of a microalgae strain. The strain grew in a culture medium, based on organic fertilizer. Theoretical analyses allowed the design of three primers based on the alignment of eight sequences obtained from NCBI, applying the Geneious Prime® 2019 and V1.3 and Oligo Calculator version 3.2. programs. The *in silico* restriction patterns was obtained with the NEBcutter v2.0 program. Experimental analyses began with the extraction of the DNA using the TENS protocol, then PCR amplification using PM-016F/PM-016R and PM-001F/PM-016R primers of 18S *rDNA* and finally the product was digested with *BbvCI* and *Eco53kI*; *BstUI*, *RsaI* and *MspI* enzymes. The DNA concentration extraction reached 3200 ng μl^{-1} and a purity of 2.0. The PCR amplified two products: 950 bp and 1400 bp, which brought us closer to identifying the microalgae. The *in silico* modeling and experimental restriction patterns showed similar fragments. In this way, the efficient response of restriction enzymes was demonstrated by confirming that the PM013 strain corresponds to the *Tetraselmis* genus. This method can be considered as a fast and safe alternative to identify wild microalgae in a basic molecular biology laboratory.

Keywords:

1. Introduction

Tetraselmis (Class Chlorodendrophyceae) is a genus of microalgae that has been considered an industrial source, due to the content of proteins, vitamins, lipids, carbohydrates, fatty acids and bioactive compounds. *Tetraselmis* can influence the growth and survival of fish and crustaceans because of its antioxidant properties [1]. It also accumulates many attractive oils and starches for the production of biofuels [2].

Morphologically, *Tetraselmis* is an ovoid green cell covered by a solid wall called teak. It has a single large chloroplast and four identical flagella in two different pairs; they can form colonies during some stage of their life cycle

[3]. Species of this genus have been found in marine and freshwater ecosystems worldwide [4]. These morphological characteristics observed through light and electron microscopy have been the basis for taxonomic studies of microalgae [5, 6].

During their life cycle, the genus *Tetraselmis*, acquires complex cellular characteristics that lead to confusion between taxa, but it has been insufficient for its identification [7-9]. Important results improved since the 90's, when the taxonomy of algae introduced the application of phylogenetic analyzes with molecular markers [5, 10].

Several molecular markers have been utilized to identify

microalgae such as: *rcbL*, *ITS1*, *ITS2*, *18S rDNA*, *23S rDNA* [6, 11, 12]. The *18S rDNA* gene is the most widely used, because its sequence has highly conserved flanking regions and repetitive arrangements which establish the phylogenetic relationship within distant organisms and between lineages and Phylum [4,13,10,14]. In this way, *Chlorella sorokiniana* [15], *Tetraselmis indica* [4], *Dunaliella salina* [8] and diatoms [16] were identified.

Commonly, the identification of genes is realized with DNA extraction, the amplification of a specific product and its sequencing [17, 18, 14]. However, mapping genes, using restriction enzymes, generates fast and reliable results instead of sequencing [19]. Bioinformatics programs can predict restriction patterns through *in silico* modeling, which is based on the sequence of the molecular marker and the relative location of the restriction sites for different enzymes [20-21]. *In silico* modeling provides the size of the DNA fragments and can subsequently be validated experimentally [22, 19].

This work is being proposed to compare *in silico* modeling with experimental restriction patterns based on the *18S rDNA* gene for the identification of a microalgae strain isolated from the Santa Elena Peninsula - Ecuador, morphologically identified as *Tetraselmis*.

2. Materials and Methods

2.1. Microalgae Strain and Cultivation

The sample was obtained from the Santa Elena Peninsula, Ecuador (2 ° 11'36.7 "S 80 ° 59'10.0" W) and registered with the code PM013 corresponding to the project INCYT-PNF-2017M3121. The strain was cultivated following a previous methodology [23]. The culture medium was prepared in 200 ml bottles based on an organic fertilizer (2% NPK) (Marchfolm), containing sodium metasilicate as an inorganic source and enriched with vitamins (1 ml l⁻¹). The cultures were installed in triplicate for six days under the following conditions: temperature of 24 ± 0.5°C, a photon flux density of 60 µmol m⁻² s⁻¹, provided by white light fluorescent lamps (Sylvania 18W), photoperiod 12/12 hours (light/dark), pH 7, shake manually twice daily, following a previous protocol [24]. Biomass was obtained by centrifugation at 13000 rpm.

2.2. Theoretical Molecular Analysis

2.2.1. Alignments and Primers Design

Eight sequences of the *18S rDNA* gene were selected from GenBank and aligned with the Clustal method using the ClustalW Alignment v2.0.12 program to obtain a conserved region and from this alignment, three *Tetraselmis* specific primers were designed: PM-001F *Forwards* (5'-GAGTGTTCAAAGCAAGCCTACG -3'), PM-016F *Forwards* (5'- AGAAACGGCTACCACATCCA -3') and PM-016R *Reverse* (5'-TGTACAAAGGGCAGGGACGTAATC-3'), using Geneious Prime® 2019 software. v1.3 and Oligo Calculator v3.27 [25].

2.2.2. In Silico Modeling of the 18S rDNA Gene

To obtain the *Tetraselmis in silico* restriction pattern, we used the consensus sequence from the *18S rDNA* and it was analyzed by the BioEdit program Map restriction and NEBcutter v2.0. The enzymes *BstUI*, *MspI*, *RsaI*, *BbvCI* and *Eco53kI* were selected.

2.3. Experimental Molecular Analysis

2.3.1. Total DNA Extraction

The extraction of the total DNA was carried out using a modifying TENS protocol [26]. In an eppendorf tube, 50 mg of *Tetraselmis*, 600 µl of TENS buffer and 2 µl of proteinase K were added and the solution was incubated at 37°C for 15 min. Then 200 µl of phenol and 200 µl of chloroform: isoamyl alcohol (24:1) was added and centrifuged at 10000 rpm for 10 minutes, the supernatant was transferred to a new tube containing 500 µl of 95% ethanol frozen, the sample was left to rest at -20°C for 30 min and subsequently centrifuged. The pellet was washed with 800 µl of 75% ethanol frozen and finally resuspended in 100 µl of UltraPure™ Water. The extracted DNA went through a purification process with sodium acetate (3M, pH 5.2) [27]. Then it was measured by spectrophotometry at an A₂₆₀/A₂₈₀ wavelength with a 2000cc Nanodrop.

2.3.2. PCR Amplification

PCR experiments were carried out using primers PM-016F and PM-016R; other with PM-001F and PM-016R. The PCR reactions were performed in a total volume of 15 µl, containing 25ng of genomic DNA, 0.125 mM of each primer, 0.2 mM of dNTPs, 3mM of MgCl₂, 1 U Taq DNA Polymerase and Buffer 1X (All products from Invitrogen™) [28]. The PCR conditions were as follows: an initial denaturation of the template DNA at 94°C for 10 min, followed by 35 cycles of 94°C for 40s, 61°C for 30s and 72°C for 1 min 30s, and finished with a final extension at 72°C for 5 minutes using a thermal cycler (Bio-Rad T100). The PCR fragments were separated by 1% agarose gel electrophoresis with 0.5X GelRed (Biotium).

2.3.3. Digestion of the PCR Product

The PCR product was digested with the enzymes *BbvCI* and *Eco53kI* for the first reaction; *BstUI*, *MspI* and *RsaI* for the second. All enzymes were from Invitrogen™ and the performance was according to a previous protocol [19]. 10 µl of the PCR product was used for the enzyme reaction (1U/enzyme) in a final volume of 50 µl. The sample was incubated at 37°C for 2 hours. The product was run on a 1% agarose gel electrophoresis, with Sybr® Green staining and visualized in an UV Transilluminator.

3. Results

3.1. Sequence Alignment and Primer Design

In order to determine the conserved region of the *18S rDNA* gene, we selected eight pre-established sequences related to *Tetraselmis*, and we analyzed the genetic

relationship. The alignment showed that the species share a high percentage of similarity (99%) (Table 1). The conserved region was evident from 380 bp to 1680 bp.

Based on the conserved region, we designed three primers

(PM-016F, PM-016R and PM-004F) to identify the *Tetraselmis* genus. The Geneious Prime® 2019 program V1.3 and Oligo Calculator v3.27 provided the optimal characteristics of the primers (Table 2).

Table 1. Sequences of the genus *Tetraselmis* from GenBank used in this study.

Sequences	N° (bp)	Similarity data (%)	N° Access (Genbank)
<i>Tetraselmis</i> sp.	1674	100	GQ917221.1
<i>Tetraselmis chuii</i>	1751	99.94	JN903999.1
<i>Tetraselmis</i> sp.	1701	100	AJ431370.2
<i>Tetraselmis carteriformis</i>	1641	99.24	FJ559384.1
<i>Tetraselmis striata</i>	1674	99.26	JQ315739.1
<i>Tetraselmis suecica</i>	1674	100	JF489949.1
<i>Tetraselmis subcordiformis</i>	1719	99.01	KU561107.1
<i>Tetraselmis marina</i>	1669	99.77	KY045847.1

Table 2. Characteristics of the primers designed for the amplification of the *Tetraselmis* 18S rDNA gene.

Reaction	Primer	Sequences (5'→3')	Length (bp)	G+C (%)	Tm ¹ (°C)	Ta ² (°C)	Product length (bp)
1	PM-016F	AGAAACGGCTACCACATCCA	20	50	59	61	1400
	PM-016R	TGTACAAAGGGCAGGGACGTAATC	24	50	62		
2	PM-001F	GAGTGTTCAAAGCAAGCCTACG	22	50	61	59	950
	PM-016R	TGTACAAAGGGCAGGGACGTAATC	24	50	62		

¹Tm: temperature medium theoretical

²Ta: temperature hybridization used in this study

3.2. Quality of DNA Extracted and Amplification by PCR

The quality of the extracted DNA was 3200 ng µl⁻¹ and its purity was 2.0 under the absorbance ratio A₂₆₀/A₂₈₀. Due to high concentration of DNA, it was diluted (1:25) to be used in PCR amplification.

From the theoretical analysis, the size of the first PCR product was 1400 bp with PM-016F/PM-016R and the second product was 950 bp with PM-001F/PM-016R. This would allow the amplification of much of the length of the 18S rDNA in *Tetraselmis* species. The experimental results displayed the same desired bands (Figures 1 and 2), demonstrating that the PM013 strain would correspond to the *Tetraselmis* genus.




Figure 1. PCR of 18S rDNAs amplified with PM-016F/PM-016R primers. Lanes 1 to 3: different samples of *Tetraselmis* (PM013). C-: negative control M: molecular weight marker (1kb Invitrogen™). Right arrow indicates the size of the fragment.




Figure 2. PCR of 18S rDNAs amplified with PM-001F/PM-016R primers. Lanes 1 to 3: different samples of *Tetraselmis* (PM013). M: molecular weight marker (1kb Invitrogen™). Right arrow indicates the size of the fragment.

3.3. Comparison of the in Silico Versus Experimental Restriction Pattern

The enzymes *BbvCI* and *Eco53kI* digested a PCR product of the first reaction and the enzymes *BstUI*, *RsaI* and *MspI* digested the second reaction. The *in silico* modeling predicted restriction patterns where each enzyme made more than two cuttings along it (Figure 3A and 4A).

The experimental restriction patterns showed fragments of similar sizes to the *in silico* modeling. The *BbvCI* performed two cuttings: 900 bp and 500 bp, while *Eco53kI* produced a fragment of 1000 bp and the other of 400 bp. The *BstUI* revealed three cuttings: a 700 bp fragment and another of approximately 200 bp; the 50 bp fragment was unseen. The *RsaI* enzyme produced two cuttings, the largest fragment was 900 bp and the small fragment was unseen in the 1% agarose gel. The *MspI* generated three cuttings, a 400 bp fragment and two overlapping fragments of approximately 250 bp,

while the smallest fragment was impossible to show (Figure 3B and 4B).




Figure 3. A) *In silico* modeling of the restriction pattern of the *18S rDNA* gene *Tetraselmis* digested with *BbvCI* and *Eco53kI* enzymes. B) Experimental restriction pattern of the *18S rDNA* gene digested with *BbvCI* and *Eco53kI* enzymes. C-: undigested PCR product; M: molecular weight marker (1Kb InvitrogenTM) 1% agarose gel. The arrows indicate the size of the fragments.




Figure 4. A) *In silico* modeling of the restriction pattern of the *18S rDNA* gene *Tetraselmis* digested with *BstUI*, *RsaI* and *MspI* enzymes. B) Experimental restriction pattern of the *18S rDNA* gene digested with *BstUI*, *RsaI* and *MspI* enzymes. C-: undigested PCR product; M: molecular weight marker (1Kb InvitrogenTM) 1% agarose gel. The arrows indicate the size of the fragments.

4. Discussion

We identified a *Tetraselmis* sp. microalgae strain isolated from the Santa Elena Peninsula, Ecuador, and compared it with *in silico* modeling and experimental restriction patterns based on the *18S rDNA* gene.

The sequence comparison allows us to infer that the similarity and evolution grade of a new molecule can be based on a database such as NCBI [29, 30]. Taking advantage of the existing information, we performed the alignment of eight sequences of the *18S rDNA* gene from various *Tetraselmis* species, to find the conserved region; this allowed us to obtain a more reliable result with real data. Furthermore, according to a previous work [13], sequence conservation is one of the most direct ways of estimating the precision of an alignment in computational biological models and the use of a bioinformatic program that provides us with accurate data such as Clustal W Alignment v2.0.12.

The conserved region of the *Tetraselmis 18S rDNA* gene facilitated the design of the three primers (PM-016F, PM-

016R and PM-001F) used in this study. According to a previous work, primer design should start with sequence alignment to ensure specificity [34].

The size of the *18S rDNA* gene in *Tetraselmis* has been described previously, and it oscillates around 2500 bp [32]. However, other studies found PCR products between 1000 bp and 1500 bp [33, 34, 28]. This evidence agrees with our results, because we amplified two products, one of 950 bp and the other of 1400 bp, this fact allowed us to take steps toward the molecular identification of the PM013 strain as a *Tetraselmis* sp.

A strong PCR amplification also depends on the quality of the template DNA. The TENS protocol has been efficient for the extraction of total DNA in microorganisms, especially in microalgae, because it contributes to the proteins denaturation of the cell wall that causes a total lysis of the cell [35], high concentration and purity [31, 26, 34]. We obtained DNA of high quality in concentration and purity which we recommend for molecular studies of microalgae.

In this era of new knowledge, the molecular identification

has contributed to deciphering species at any stage of their life cycle [9, 17, 16]. The PCR product has been generally sequenced and evaluated using bioinformatic programs that ensure the functional or phylogenetic relationship [13, 16, 14]. However, the benefit of using restriction enzymes could be an alternative technique based on the PCR product and is faster and reliable [21, 28, 19]. For example, Olmos, *et al.* (2000) implemented this technique in the molecular identification of *Dunaliella* sp [8]; while Chin, *et al.* (2018) and Pillacela, *et al.* (2020) performed an *in silico* analysis to obtain the restriction pattern before being experimentally evaluated *Pyrodictum bahamense* [34] and *Haematococcus* sp [19], respectively.

In our study, we predicted the *in silico* restriction patterns for two PCR products, and then we digested experimentally using the enzymes *BbvCI* and *Eco53kI*; *BstUI*, *RsaI*, and *MspI*. The results were satisfactory, the experimental restriction patterns were highly related to the *in silico* modeling, although small fragments (<100 bp) were unseen because they correspond to the ends of the sequence, left by the restriction enzymes cuttings, and are generally imperceptible and lack biological validity. The same variations were presented by Pillacela, *et al.* (2020) and Cienfuegos, *et al.* (2008) [19, 39].

Currently, *in silico* modeling is important in molecular biological studies [36, 18] because they predict reliable results and save time and money in the experimental stage [37, 38]. The high similarity between the restriction patterns obtained *in silico* and experimentally demonstrated the veracity of the results. This suggests the PM013 strain, isolated from the Santa Elena Peninsula Ecuador, is related to *Tetraselmis* sp. In addition, laboratories can work in basic molecular biology to identify organisms at the genus level.

5. Conclusions

Theoretical analyses allowed to find the conserved region of the *Tetraselmis* 18S rDNA gene, thus it facilitated the design of primers PM-016F/PM-016R and PM-001F/PM-016R, which provided PCR products of 1400 bp and 950 bp, respectively. We could show similar fragments between the *in silico* modeling and experimental restriction patterns after the enzymes *BbvCI* and *Eco53kI*; *BstUI*, *RsaI* and *MspI* digested each product, which confirms that the PM013 strain corresponds to the *Tetraselmis* sp. Those results were supported by high quality DNA obtained with the TENS protocol, which allows us to suggest its usefulness in molecular biology studies of the *Tetraselmis*. Finally, we infer that bioinformatics applications and *in silico* modeling are essential before experimental tests of biological processes, in this work they allowed us a fast and effective response of restriction enzymes when carrying out molecular identification in a basic laboratory where sequencing work is difficult.

Acknowledgements

This research was supported by the Center for

Advanced Studies of Arid Zones (CEAZA-Chile) through Dr. Claudio Álvarez and financed by the INCYT-PNF-2017M3112 Projects (Ecuador); UPSE-CUP: 91870000.0000.384095 (Ecuador), with the permission of the Ministry of the Environment: MAE-DNB-CM-CM-2018-0099. We thank the MsC. Jimmy Villón, and Blgo. Bryan Pillacella for his support in the sampling and cultivation of the microalgae.

References

- [1] Macías, P., Coello, D., & García, D. (2018). Floración de *Tetraselmis* sp., en la zona costera frente a las Provincias de Manabí y Santa Elena (Abril 2018). Guayaquil: Instituto Nacional de Pesca.
- [2] Úbeda, P., Chileh, T., Dautor, Y., García, F., & Alonso, D. (2015). Tools for microalgal biotechnology: Development of an optimized transformation method for an industrially promising microalga *Tetraselmis chuii*. *J. Appl. Phycol.*, 1 (27), 223-232.
- [3] Guiry, M., & Guiry, G. (2018). *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. Recuperado el 11 de Junio de 2019, de World-wide electronic publication, National University of Ireland, Galway: <http://www.algaebase.org>
- [4] Arora, Mani. (2016). *Tetraselmis*: an Introducción. *The Botanica*, 66. 155-175.
- [5] Juliet, B., & Lewis, J. (2007). Unravelling the algae. the past, present and future of algal systematics. London, UK: *Systematics Association special volume series*, 75.
- [6] Arora, M., Anil, A., Leliaert, F., Delany, J., & Mesbahi, E. (2013). *Tetraselmis indica* (Chlorodendrophyceae, Chlorophyta), a new species isolated from salt pans in Goa, India. *European Journal of Phycology*, 48 (1), 61–78.
- [7] Lee, H. & Hur, S., (2009). Genetic Relationships among Multiple Strains of the Genus *Tetraselmis* Based on Partial 18S rDNA Sequences. *Algae*, 24 (4), 205-212.
- [8] Olmos, J., Paniagua, J., & Contreras, R. (2000). Molecular identification of *Dunaliella* sp. utilizing the 18S rDNA gene. *Letters in Applied Microbiology*, 80–84.
- [9] Zou, S., Fei, C., Wang, C., Gao, Z., Bao, Y., He, M., & Wang, C. (2016). How DNA barcoding can be more effective in microalgae identification: a case of cryptic diversity revelation in *Scenedesmus* (Chlorophyceae). *Scientific reports*, 6, 36822. 10.1038/srep36822.
- [10] Tragin, M., Lopes dos Santos, A., Christen, R., & Vaultot, D. (2016). Diversity and ecology of green microalgae in marine systems: an overview based on 18S rRNA gene sequences. *Perspectives in Phycology*, 3 (3), 141–154.
- [11] Kusumawati, L., Wahyudi, R., Pinontoan, R., Gorreti, M., & Panggabean, L. (2015). Sequence Analysis of 18S DNA of *Melosira* sp., *Dunaliella* sp., *Isochrysis* sp. and *Porphyridium* sp. *The 3rd International Conference on Biological Science* 2013, 2, 592–597.
- [12] Turmel, M., De Cambiaire, J., Otis, C., & Lemieux, C. (2016). Distinctive architecture of the chloroplast genome in the chlorodendrophycean green algae *Scherffelia dubia* and *Tetraselmis* sp. CCMP 881. *PLoS ONE*, 11 (2), 1-22.

- [13] Chatzou, M., Cedrik, M., Jia Ming, C., Carsten, K., Giovanni, B., Ionas, E., & Cedric, N. (2016). Multiple sequence alignment modeling: methods and applications. *Oxford University*, 17 (6), 1009–1023.
- [14] Gunapati, O., A. Thadoi., K. Ojit Singh., O. Avijeet Singh., Th. Indrama., Laxmipriya Kojiam., Wangkhem, Indira., Chungkham, Silvia., A. Sharma., Romi, Khangembam., Minerva, Shamjetshabam., Th. Bidyababy., K. Sarabati., O.N. Tiwari & G.D. Sharma (2015). Molecular characterization of selected diazotrophic cyanobacteria of North East India for phylogenetic study *Int. J. of Adv. Res.* 3 (8), 261-271.
- [15] Flores, F., Maldonado, S., & Ortiz, J. (2016). Identificación molecular de microalgas clorófitas del Ecuador. *Research Gate*, 1-2.
- [16] Benoiston, A.-S., Ibarbalz, F., Bittner, L., Guidi, L., Jahn, O., Dutkiewicz, S., & Bowler, C. (2017). The evolution of diatoms and their biogeochemical functions. *Phil. Trans. R. Soc. B*, 372 (20160397), 2-10.
- [17] Bérard, A., Dorigo, U., Humbert, J., & Martin, F. (2005). Microalgae community structure analysis based on 18S rDNA amplification from DNA extracted directly from soil as a potential soil bioindicator. *Agronomy for Sustainable Development, Springer Verlag/EDP Sciences/INRA*, 25 (2), 285-291.
- [18] Galarza, J., Gimpel, J., Rojas, V., Arredondo-Vega, B., & Henríquez, V. (2018). Over-accumulation of astaxanthin in *Haematococcus pluvialis* through chloroplast genetic engineering. *Algal Research*, 31, 291–297.
- [19] Pillacela, B., Galarza, J., & Tufiño, C. (2020). Reliability of in silico Modeling Based on Restriction Patterns for the Study of the Carotenogenic Gene pds of *Haematococcus* sp. *International Journal of Microbiology and Biotechnology*, 5 (1), 16-21.
- [20] František, M. (2005). *Physical Mapping-Restriction Mapping* (4th edition ed.). USA: Molecular Cell Biology.
- [21] Vincze, T., Posfai, J., & Roberts, R. (2003). NEBcutter: a program to cleave DNA with restriction enzymes. *Nucleic Acids Res*, 31, 3688-3691.
- [22] Parashar, D, Srivastava R, Chauhan D, Sharma V, Maharaj Singh, Mallika Lavania, Chauhan A, & Bhatia A. (2006). Characterization of mycobacteria isolated from bovines by PRA-targeting hsp 65 gene region. *J Commun Dis*, 38 (2): 263-268.
- [23] Garduño-Solórzano, G., Rodríguez, M., Martínez, M., Quintanar, R., Lozano, C., Campos, J., & Monsalvo, A. (2011). Cultivos de microalgas del Lago de Catemaco, Veracruz. *Latinoam Biotechnol Amb Algal*, 2 (2), 67-80.
- [24] Nagaraj, S., Arulmurugan, P., Rajaram, M., Sundararaj, R., & Rengasamy, R. 2012. Enhanced production of astaxanthin at different physico-chemical parameters in the green alga *Haematococcus pluvialis* Flotow. *Phykos*, 42 (1), 59–71.
- [25] Kibbe, W. (2007). Oligo Calc:Oligonucleotide Properties Calculator. Recuperado el 6 de Junio de 2019, de <http://biotools.nubic.northwestern.edu/OligoCalc.html>.
- [26] Green, M., & Sambrook, J. (2012). *Molecular Cloning*. New York: Cold Spring Harbor Laboratory Press.
- [27] Rocha, P. (2002). Teoría y práctica para la extracción y purificación del ADN de palma de aceite. Bogotá, Colombia: Laboratorio de Marcadores Moleculares, Área de Fisiología y Mejoramiento.
- [28] Yáñez, R. (2011). Categorización taxonómica en base a marcadores moleculares de microalgas extremófilas nativas del Norte de Chile. Chile: Facultad De Ciencias; Instituto De Biología.
- [29] Morrison, D., & Ellis, J. (1997). Effects of Nucleotide Sequence Alignment on Phylogeny Estimation: A Case study of 18S rDNAs of apicomplexa. *Mol. Biol. Evol*, 14 (4), 428-441.
- [30] Molins, A., Moya, P., García, F., Reig, J., & Barreno, E. (2018). A multi-tool approach to assess microalgal diversity in lichens: isolation, Sanger sequencing, HTS and ultrastructural correlations. *The Lichenologist*, 50 (1), 123–138.
- [31] Fawley, M. & Fawley, K. (2004). A simple and rapid technique for the isolation of DNA from microalgae. *Phycological Society of America*, Volumen 40, pp. 223-225.
- [32] Cruz, J. (2012). Comparación De Genes De Arn Ribosomal 18s De Tres Cepas Del Género Euglena: *E. Pailasensis*, *E. Mutabilis Cpc293* y Una Euglena Encontrada En Un Riachuelo Ácido Del Parque Nacional Volcán Rincón De La Vieja. Cartago: Instituto Tecnológico De Costa Rica.
- [33] Baskara, A., Jayakumar, T., Ganasan, M., Mohan, N., Senthil, C., Nagaraj, S., Rengasamy, R., Manubolu, M., Sheu, J., & Chang, C. (2018). Mass cultivation of new algae *Tetraselmis* straiata BBRR1 under open raceway ponds for biodiesel and biocrude production. Tamil Nadu, India: Preprints.
- [34] Chin, W., Teoh, P., Anton, A. & Kumar, S., (2018). Molecular Characterization And Identification Of Ribosomal Dna Sequences Of A Harmful Algal Bloom Species, Pyrodinium Bahamense Var. Compressum. *Biotechnology Research Institute*, 1 (1), pp. 1-8.
- [35] Montes, J., & Pulido, M. (2012). Obtención de protocolos para el aislamiento, cultivo y extracción de ADN de *Chlorella vulgaris* Beyerinck. *Investigación y Ciencia del Gimnasio Campestre* (45), 47-54.
- [36] Galarza, J., Delgado, N., & Henríquez, V. (2016). Cisgenesis and intragenesis in microalgae: promising advancements towards sustainable metabolites production. Publisher, Springer International Publishing. *Applied Microbiology and Biotechnology*, 100 (24), 10225–10235.
- [37] Cagney, G., Amiri, S., Premawaradena, T., Lindo, M., & Emili, A. (2003). In silico proteome analysis to facilitate proteomics experiments using mass spectrometry. *Proteome Science*, 1 (1), 5.
- [38] Jiménez, J., & Chaparro, A. (2016). In silico design and functional assessment of semisynthetic genes that confer tolerance to phosphinothricin. *Revista Colombiana de Biotecnología*, XVIII (2).
- [39] Cienfuegos, A., Conn, J., Gomez, G., & Correa, M. (2008). Diseño y evaluación de metodologías basadas en PCR-RFLP de ITS2 para la identificación molecular de mosquitos *Anopheles* spp (Diptera: Culicidae) de la Costa Pacífica de Colombia. *Rev Biomed*, 19 (1), 35–44.